NOAA Technical Report NMFS 150

August 2000

Relationship between Fish Size and Otolith Length for 63 Species of Fishes from the Eastern North Pacific Ocean

James T. Harvey
Thomas R. Loughlin
Michael A. Perez
Dion S. Oxman

U.S. Department of Commerce

AA
ch.
pt.
FS
0
1

Library
NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

U.S. DEPARTMENT OF COMMERCE

NORMAN Y. MINETA
SECRETARY

National Oceanic and Atmospheric Administration

D. James Baker
Under Secretary for
Oceans and Atmosphere

National Marine Fisheries Service

Penelope D. Dalton
Assistant Administrator
for Fisheries

NOAA Technical Reports NMFS

Technical Reports of the *Fishery Bulletin*

Scientific Editor
Dr. John V. Merriner
Southeast Fisheries Science Center
National Marine Fisheries Service, NOAA
101 Pivers Island Road
Beaufort, North Carolina 02516

Editorial Committee
Dr. Andrew E. Dizon National Marine Fisheries Service
Dr. Linda L. Jones National Marine Fisheries Service
Dr. Richard D. Methot National Marine Fisheries Service
Dr. Theodore W. Pietsch University of Washington
Dr. Joseph E. Powers National Marine Fisheries Service
Dr. Tim D. Smith National Marine Fisheries Service

Managing Editor
Shelley Arenas
Scientific Publications Office
National Marine Fisheries Service, NOAA
7600 Sand Point Way N.E.
Seattle, Washington 98115-0070

The *NOAA Technical Report NMFS* (ISSN 0892-8908) series is published by the Scientific Publications Office, National Marine Fisheries Service, NOAA, 7600 Sand Point Way N.E., Seattle, WA 98115-0070.

The Secretary of Commerce has determined that the publication of this series is necessary in the transaction of the public business required by law of this Department. Use of funds for printing of this series has been approved by the Director of the Office of Management and Budget.

The *NOAA Technical Report NMFS* series of the *Fishery Bulletin* carries peer-reviewed, lengthy original research reports, taxonomic keys, species synopses, flora and fauna studies, and data-intensive reports on investigations in fishery science, engineering, and economics. The series was established in 1983 to replace two subcategories of the Technical Report series: "Special Scientific Report—Fisheries" and "Circular." Copies of the NOAA Technical Report NMFS are available free in limited numbers to government agencies, both federal and state. They are also available in exchange for other scientific and technical publications in the marine sciences.

NMFS Honolulu Library
00000000004877

NOAA Technical Report NMFS 150
A Technical Report of the *Fishery Bulletin*

Relationship between Fish Size and Otolith Length for 63 Species of Fishes from the Eastern North Pacific Ocean

James T. Harvey
Thomas R. Loughlin
Michael A. Perez
Dion S. Oxman

August 2000

NOAA
Tech
Rept
NMFS
150
C.1

U.S. Department of Commerce
Seattle, Washington

Library
NOAA, National Marine Fisheries Service
SWFC Honolulu Laboratory F/SWC2
2570 Dole Street
Honolulu, HI 96822-2396

Suggested reference
Harvey, James T., Thomas R. Loughlin, Michael A. Perez, and Dion S. Oxman. 2000. Relationship between fish size and otolith length for 63 species of fishes from the eastern North Pacific Ocean. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 150, 36 p.

Online dissemination
This report is posted online in PDF format at http://spo.nwr.noaa.gov (click on *Technical Reports* link).

Purchasing additional copies
Additional copies of this report are available for purchase in paper copy or microfiche from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161; 1-800-553-NTIS; http://www.ntis.gov.

Copyright law
Although the contents of the Technical Reports have not been copyrighted and may be reprinted entirely, reference to source is appreciated.

Proprietary products
The National Marine Fisheries Service (NMFS) does not approve, recommend, or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

Relationship between Fish Size and Otolith Length for 63 Species of Fishes from the Eastern North Pacific Ocean

JAMES T. HARVEY*, THOMAS R. LOUGHLIN, MICHAEL A. PEREZ, and DION S. OXMAN**

National Marine Mammal Laboratory
Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
7600 Sand Point Way N.E.
Seattle, Washington 98115-0070

ABSTRACT

Otoliths commonly are used to determine the taxon, age, and size of fishes. This information is useful for population management, predator-prey studies, and archaeological research. The relationship between the length of a fish and the length of its otoliths remains unknown for many species of marine fishes in the Pacific Ocean. Therefore, the relationships between fish length and fish weight, and between otolith length and fish length, were developed for 63 species of fishes caught in the eastern North Pacific Ocean. We also summarized similar relationships for 46 eastern North Pacific fish species reported in the literature. The relationship between fish length and otolith length was linear, and most of the variability was explained by a simple least-squares regression ($r^2 > 0.700$ for 45 of 63 species). The relationship between otolith length and fish length was not significantly different between left and right otoliths for all but one fish species. Images of otoliths from 77 taxa are included to assist in the identification of species.

Introduction

All bony fishes (Osteichthyes) have three pairs of otoliths (earbones or earstones): the sagittae, asteriscus, and lapillus. These otoliths are composed of calcium carbonate in the form of aragonite, in a protein matrix. They are contained within membranous labyrinths in paired otic capsules on either side of the skull. The sagittae are the largest pair of otoliths in most bony fishes; however, in minnows (Cypriniformes) and catfish (Siluriformes) the asterici are the largest (Hecht, 1977). Fisheries biologists have used sagittae to determine age and growth of fishes because of the large size and distinct growth rings of sagittae (Chilton and Beamish, 1982; Boehlert, 1985; Summerfelt and Hall, 1987).

Because otoliths are dense they can withstand some degree of dissolution, and often species can be recognized by the distinctive morphology of the sagittae (Morrow, 1979; Smale et al., 1995). Paleontologists have identified otoliths in middens (Fitch, 1969), oceanographers have determined species of fishes from otoliths in sediments (Fitch, 1964, 1968), and prey have been identified using otoliths collected from stomachs of piscivorous fishes (Trippel and Beamish, 1987), marine birds (Ainley et al., 1981), and marine mammals (Fitch and Brownell, 1968; Treacy and Crawford, 1981). Fishes eaten by pinnipeds also were identified using otoliths found in feces (Bailey and Ainley, 1982; Brown and Mate, 1983; Antonelis et al., 1984; Harvey, 1987).

Trout (1954) and Templemann and Squires (1956) were among the first to demonstrate a significant positive relationship between otolith size and fish size of Barents Sea cod (*Boreogadus saida*) and haddock (*Melanogrammus aeglefinus*). Otolith length also has been correlated with fish weight (Casteel, 1976). Since these early studies, relationships between otolith length and fish length have been determined for some species, including North Pacific gadids (Frost and Lowry, 1981), rockfishes (*Sebastes* spp.; Wyllie Echeverria, 1987), and several fishes off Baja California, Mexico (Gamboa, 1991).

For most species, the relationship between otolith length and fish length can be described by a simple linear regression. For North Pacific gadids, this relation-

* Present address: Moss Landing Marine Laboratories, 8272 Moss Landing Road, Moss Landing, CA 95039-9647.

** Present address: University of Alaska, Fairbanks, Juneau Center, School of Fisheries and Ocean Sciences, 11120 Glacier Hwy, Juneau, Alaska 99801.

ship has been best described by two linear regressions with an inflection point (Frost and Lowry, 1981). Left and right sagittae also may differ in size within a rockfish species (Wyllie Echeverria, 1987), and sometimes otolith size is different among stocks of fishes, such as Atlantic herring (*Clupea harengus*; Messieh, 1972).

The objective of this study was to compile information regarding the relationship between otolith length and fish length for fish species of the eastern North Pacific Ocean using original collections and published and unpublished literature. These data may be used by researchers studying archaeology and food habits of piscivores to determine the size of fishes from the length of recovered otoliths. We also wanted to provide images of most fish otoliths, for which we had regressions, to be used as an aid for identifying species of fish.

Methods

Fishes were collected throughout the eastern North Pacific Ocean (e.g. Bering Sea, Gulf of Alaska, and off Washington, Oregon, and California) using bottom and midwater trawls, beach seines, gill nets, and hook-and-line gear. Fish were weighed to the nearest 0.1 g on a Mettler balance (<600 g) and spring or pan scale (>600 g). Standard length (SL; most anterior point to the base of hypural plate at caudal flexion) or fork length (FL; most anterior point to the base of the fork in the caudal fin) was measured to the nearest millimeter.

Sagittae were removed, cleaned, and stored dry in vials. Lengths of sagittae were determined using handheld vernier calipers under a dissecting microscope. Sagittal otolith length was recorded as the greatest distance measured from the anterior rostrum to the posterior edge, parallel to the sulcus.

The relationship between otolith length and fish length (SL or FL) was determined using a least-squares linear regression. The appropriateness of the linear model was determined by plotting the residuals against the independent variable. Differences between regression coefficients for the relationship of fish length and the lengths of left and right otoliths were tested using t tests. When the equations for left and right otoliths did not differ statistically, one right or left otolith was selected randomly from each individual and a single linear regression reported for each fish species. The significance of the linear regression was tested using an analysis of variance (ANOVA). Relationships between otolith length and fish length for additional species were obtained from published and unpublished sources. Relationships between fish length and fish weight were determined using a least-squares regression of the log of fish weight and length (Ricker, 1975). Although transformation back to arithmetic units may result in underestimating weight, these errors are usually small (Saila et al., 1988).

Results

Forty-six relationships between fish and otolith size previously reported in the literature involved various measures of fish length—fork length, total length (defined as the distance from the most anterior point to the most posterior point), and standard length measured in millimeters or centimeters—and otolith length (Table 1). Many of the published regressions of fish length to otolith length were developed for species common in food habit studies of marine mammals (Frost and Lowry, 1981; Brown and Mate, 1983) or species that were commercially important (Spratt, 1975; Boehlert and Yoklavich, 1984; Wyllie Echeverria, 1987).

Sixty-three species of fishes were collected in connection with the current study (Table 2). Most relationships between weight and length were described by a traditional allometric equation, where weight of a fish is approximately equal to length to the third power (Table 2). Linearized forms of this power relationship explained >90% of the variability in 43 of 60 cases. For three species, no weight data were collected. For 17 additional species, the sample size was less than 20; therefore, these weight/length relationships should be used with caution.

Generally the relationship between fish length and otolith length was linear, and most of the variability was explained by the regression (r^2>0.700 in 45 of 63 cases; Table 2). All relationships except one were significant (P<0.05). There was no significant relationship between otolith length and fish length for *Trachurus symmetricus* (Table 2). Regression coefficients of otolith length to fish length were not significantly different for left and right otoliths, except for the wattled eelpout (*Lycodes palearis*; P<0.05), however, the analysis was probably influenced by the small sample size (n=12). Size of fish should not be predicted from otolith size or fish weight for measurements outside the range used for the regressions.

To assist in the identification of recovered otoliths, we provide images of fish otoliths (Fig. 1) for most species sampled. These otoliths are listed according to taxonomic relationships (Robins et al., 1991). We chose otoliths that were representative of the species, and presented multiple images of otoliths from species where the otolith morphology changed with size.

Discussion

Otoliths have been used to identify fish species eaten by marine predators (Fitch and Brownell, 1968; Pitcher, 1980;

Table 1

Published relationships of fish otolith length (or mass) and fish length (or mass) for fishes in the eastern North Pacific. NA indicates data not available, and the codes for the variables (x and y) are defined at the bottom of the table.

Species	Regression	Variables	r^2	n	Location	Reference
Boreogadus saida (Arctic cod)	$y = 2.20x + 1.59$	b	0.98	202	Beaufort & Chukchi Seas	Frost and Lowry (1981)
Citharichthys sordidus (Pacific sanddab)	$y = 0.02x + 1.02$	a	0.93	46	Oregon	Brown and Mate (1983)
Eleginus gracilis (saffron cod)	$y = 2.32x - 4.84$	b, c	0.96	110	Bering & Chukchi Seas	Frost and Lowry (1981)
	$y = 1.74x - 0.09$	b, d	0.93	36		
Engraulis mordax (northern anchovy)	$y = 33.22x - 8.49$	e	0.88	121	California	Spratt (1975)
Errex zachirus (rex sole)	$y = 0.02x + 1.02$	a	0.98	78	Oregon	Brown & Mate (1983)
Hippoglossus stenolepis (Pacific halibut)	$\ln(y) = 1.31 \ln x - 1.32$	b	NA	2,503	North Pacific	Southward (1962)
	$\ln(y) = 1.97 \ln x + 6.00$		0.90	128	North Pacific	Southward and Chapman (1965)
Leptocottus armatus (staghorn sculpin)	$y = 0.03x + 1.31$	a	0.98	14	Oregon	Brown and Mate (1983)
Lyopsetta exilis (slender sole)	$y = 0.02x + 0.30$	a	0.98	47	Oregon	Brown and Mate (1983)
Microstomus pacificus (dover sole)	$y = 0.02x + 0.62$	a	0.97	45	Oregon	Brown and Mate (1983)
Oncorhynchus keta (chum salmon)	$\log_{10} y = 1.23 + 3.21 \log_{10} x$	f	0.99	43	North Pacific	Casteel (1974)
Oncorhynchus kisutch (coho salmon)	$\log_{10} y = 0.89 + 5.93 \log_{10} x$	f	0.99	19	North Pacific	Casteel (1974)
Oncorhynchus mykiss (rainbow trout)	$y = 0.006x + 2.16$	a	0.97	121	British Columbia, Washington, Oregon	McKern et al. (1974)
Oncorhynchus nerka (sockeye salmon)	$\log_{10} y = 0.29 + 4.13 \log_{10} x$	f	0.95	86	North Pacific	Casteel (1974)
Oncorhynchus tshawytscha (chinook salmon)	$\log_{10} y = 0.59 + 4.15 \log_{10} x$	f	0.99	53	North Pacific	Casteel (1974)
Pleuronectes vetulus (English sole)	$y = 0.03x + 0.53$	a	0.99	81	Oregon	Brown and Mate (1983)
Sebastes auriculatus (brown rockfish)	$y = 33.16x - 53.03$	g	0.94	78	California	Wyllie Echeverria (1987)
Sebastes aurora (aurora rockfish)	$y = 19.91x + 15.12$	g	0.61	71	California	Wyllie Echeverria (1987)
Sebastes carnatus (gopher rockfish)	$y = 30.57x - 39.37$	g	0.89	203	California	Wyllie Echeverria (1987)
Sebastes caurinus (copper rockfish)	$y = 30.23x + 5.10$	g	0.82	132	California	Wyllie Echeverria (1987)
Sebastes chlorostictus (greenspotted rockfish)	$y = 24.11x - 18.54$	g	0.95	174	California	Wyllie Echeverria (1987)
Sebastes chrysomelas (black-and-yellow rockfish)	$y = 28.61x - 21.78$	g	0.84	166	California	Wyllie Echeverria (1987)
Sebastes constellatus (starry rockfish)	$y = 25.27x - 37.48$	g	0.96	99	California	Wyllie Echeverria (1987)
Sebastes crameri (darkblotched rockfish)	$y = 28.10x - 27.10$	g	0.94	89	California	Wyllie Echeverria (1987)
Sebastes diploproa (splitnose rockfish)	$y = 22.64x - 12.85$	g	0.96	78	California	Wyllie Echeverria (1987)
Sebastes elongatus (greenstriped rockfish)	$y = 24.02x - 13.56$	g	0.95	98	California	Wyllie Echeverria (1987)
Sebastes entomelas (widow rockfish)	$y = 33.11x - 6.89$	g	0.81	106	California	Wyllie Echeverria (1987)
Sebastes flavidus (yellowtail rockfish)	$y = 26.51x - 10.95$	g	0.88	184	California	Wyllie Echeverria (1987)
Sebastes goodei (chilipepper)	$y = 29.13x - 58.00$	g	0.96	78	California	Wyllie Echeverria (1987)
Sebastes hopkinsi (squarespot rockfish)	$y = 28.87x - 30.55$	g	0.92	59	California	Wyllie Echeverria (1987)
Sebastes jordani (shortbelly rockfish)	$y = 22.10x - 2.31$	g	0.97	183	California	Wyllie Echeverria (1987)

Continued

Table 1 (continued)

Species	Regression	Variables	r^2	n	Location	Reference
Sebastes levis (cowcod)	$y = 47.46x - 170.11$	g	0.95	29	California	Wyllie Echeverria (1987)
Sebastes maliger (quillback rockfish)	$y = 29.97x - 53.11$	g	0.86	34	California	Wyllie Echeverria (1987)
Sebastes melanops (black rockfish)	$y = 30.56x - 48.22$	g	0.86	209	California	Wyllie Echeverria (1987)
Sebastes melanostomus (blackgill rockfish)	$y = 30.56x - 47.07$	g	0.86	80	California	Wyllie Echeverria (1987)
Sebastes miniatus (vermilion rockfish)	$y = 29.36x - 56.74$	g	0.93	99	California	Wyllie Echeverria (1987)
Sebastes mystinus (blue rockfish)	$y = 29.77x - 18.18$	g	0.83	204	California	Wyllie Echeverria (1987)
Sebastes nebulosus (China rockfish)	$y = 25.18x + 32.97$	g	0.79	48	California	Wyllie Echeverria (1987)
Sebastes ovalis (speckled rockfish)	$y = 33.56x - 53.47$	g	0.91	84	California	Wyllie Echeverria (1987)
Sebastes paucispinis (bocaccio)	$y = 41.09x - 77.09$	g	0.82	86	California	Wyllie Echeverria (1987)
Sebastes pinniger (canary rockfish)	$y = 29.41x - 85.11$	g	0.92	173	California	Wyllie Echeverria (1987)
Sebastes rosaceus (rosy rockfish)	$y = 22.53x - 83.48$	g	0.81	147	California	Wyllie Echeverria (1987)
Sebastes ruberrimus (yelloweye rockfish)	$y = 31.33x - 76.23$	g	0.92	102	California	Wyllie Echeverria (1987)
Sebastes saxicola (stripetail rockfish)	$y = 23.40x - 32.77$	g	0.95	102	California	Wyllie Echeverria (1987)
Sebastes semicinctus (halfbanded rockfish)	$y = 25.27x - 19.18$	g	0.85	31	California	Wyllie Echeverria (1987)
Sebastes serranoides (olive rockfish)	$y = 29.35x - 51.01$	g	0.93	130	California	Wyllie Echeverria (1987)
Theragra chalcogramma (walleye pollock)	$y = 3.18x - 9.77$	b, h	0.97	98	Bering Sea	Frost and Lowry (1981)
	$y = 2.25x - 0.51$	b, l	0.98	158		

[a] x = fish standard length (mm), y = otolith length (mm)
[b] x = otolith length (mm), fish fork length (cm)
[c] otolith length > 8.5 mm
[d] otolith length < 8.5 mm
[e] x = otolith length (mm), fish standard length (mm)
[f] x = otolith length (mm), x = fish weight (g)
[g] x = otolith length (mm), x = fish total length (mm)
[h] otolith length > 10.0 mm
[i] otolith length < 10.0 mm

Table 2

Relationships of fish weight (WT) in grams to fish standard length (SL) in centimeters, and standard length to length of otolith (OL) in millimeters. For each equation, the number of fish or otoliths measured (N), coefficient of determination (r^2), standard error of the regression coefficient (SE_β), and range of fish lengths is given. Species and common names from Robins et al. (1991).

		Fish weight/fish length			Fish length/otolith length			
Species (common name)	Location[1]	Equation	N	r^2	Equation	N	r^2	Range
Allosmerus elongatus (whitebait smelt)	OR	$WT = 0.0063\ SL^{3.233}$	25	0.893	SL = 2.11 (OL) + 3.02	23	0.838	7.9–9.7
Alosa sapidissima (American shad)	OR	$WT = 0.0135\ SL^{3.046}$	7	0.997	SL = 11.46 (OL) – 11.08 $SE_\beta = 0.680$	14	0.960	8.4–37.1
Ammodytes hexapterus (Pacific sand lance)	CA AK	$WT = 0.0063\ SL^{2.790}$	10	0.913	SL = 4.06 (OL) + 2.01 $SE_\beta = 1.643$	10	0.433	9.3–13.6
Anoplopoma fimbria (sablefish)	OR	$WT = 0.0163\ SL^{2.902}$	74	0.993	SL = 5.28 (OL) + 1.62 $SE_\beta = 0.120$	94	0.955	9.3–46.7
Atheresthes stomias (arrowtooth flounder)	OR	$WT = 0.0093\ FL^{2.999}$	101	0.961	FL = 4.75 (OL) – 2.96	84	0.925	31.1–37.5
Atherinops affinis (topsmelt)	CA	$WT = 0.1698\ SL^{1.733}$	14	0.429	SL = 3.72 (OL) + 0.55 $SE_\beta = 0.325$	18	0.891	4.1–14.1
Atherinopsis californiensis (jacksmelt)	CA	$WT = 0.0049\ SL^{3.228}$	12	0.968	SL = 4.85 (OL) – 2.46 $SE_\beta = 0.279$	18	0.950	13.0–32.6
Bathymaster signatus (searcher)	WA	$WT = 0.0038\ SL^{3.256}$	44	0.991	SL = 3.48 (OL) + 1.90	43	0.883	11.8–31.5
Chilara taylori (spotted cusk-eel)	CA	$WT = 0.0004\ SL^{3.761}$	22	0.964	SL = 2.51 (OL) + 2.15 $SE_\beta = 0.294$	29	0.730	11.4–25.2
Chitonotus pugetensis (roughback sculpin)	WA	$WT = 0.0217\ SL^{2.871}$	11	0.960	SL = 3.37 (OL) – 4.52 $SE_\beta = 0.460$	11	0.857	5.5–11.8
Citharichthys sordidus (Pacific sanddab)	WA, OR	$WT = 0.0352\ SL^{2.710}$	60	0.851	SL = 2.87 (OL) + 3.29 $SE_\beta = 0.229$	61	0.727	3.3–25.5
Clupea pallasi (Pacific herring)	WA, OR	$WT = 0.0044\ SL^{3.398}$	83	0.976	SL = 5.24 (OL) – 1.85 $SE_\beta = 0.166$	82	0.934	5.1–22.7
Coryphaenoides acrolepis (Pacific rattail)	OR	$WT = 0.0016\ SL^{3.209}$	10	0.921	SL = 3.44 (OL) – 3.23 $SE_\beta = 0.368$	10	0.926	19.0–36.0
Cymatogaster aggregata (shiner perch)	OR, CA	$WT = 0.0100\ SL^{3.515}$	85	0.979	SL = 1.74 (OL) – 0.52 $SE_\beta = 0.053$	90	0.925	4.8–12.2
Dasycottus setiger (spinyhead sculpin)	GA	No data available			FL = 3.11 (OL) – 7.03 $SE_\beta = 0.368$	49	0.655	8.0–33.0
Eleginus gracilis (saffron cod)	GA	$WT = 0.0039\ SL^{3.292}$	13	0.990	SL = 1.89 (OL) – 2.76 $SE_\beta = 0.058$	46	0.960	9.2–20.9
Embiotoca jacksoni (black perch)	CA	$WT = 0.0282\ SL^{3.148}$	19	0.992	SL = 2.45 (OL) – 2.61 $SE_\beta = 0.082$	52	0.947	5.2–18.4
Embiotoca lateralis (striped seaperch)	OR, CA	$WT = 0.0329\ SL^{3.043}$	25	0.998	SL = 2.90 (OL) – 5.68 $SE_\beta = 0.061$	25	0.990	6.6–26.4
Engraulis mordax (northern anchovy)	CA	$WT = 0.0485\ SL^{2.413}$	34	0.807	SL = 2.28 (OL) + 0.85 $SE_\beta = 0.206$	56	0.694	3.6–14.4
Eopsetta exilis (slender sole)	GA	$WT = 0.0058\ SL^{3.293}$	50	0.974	SL = 3.37 (OL) + 1.08 $SE_\beta = 0.265$	50	0.771	8.0–20.5

Continued

Table 2 (continued)

Species (common name)	Location[1]	Fish weight/fish length			Fish length/otolith length			Range
		Equation	N	r^2	Equation	N	r^2	
Eopsetta jordani (petrale sole)	OR, CA	$WT = 0.0086\ SL^{3.231}$	17	0.986	$SL = 4.85\ (OL) - 4.81$ $SE_\beta = 0.468$	20	0.857	13.7–37.0
Errex zachirus (rex sole)	WA, OR	$WT = 0.0238\ SL^{2.692}$	67	0.932	$SL = 4.80\ (OL) - 2.50$ $SE_\beta = 0.226$	70	0.869	12.0–29.7
Gadus macrocephalus (Pacific cod)	BS	No data available			$FL = 4.51\ (OL) - 22.97$	110	0.883	10.0–106.0
Genyonemus lineatus (white croaker)	CA	$WT = 0.0550\ SL^{2.700}$	40	0.767	$SL = 1.52\ (OL) + 4.66$ $SE_\beta = 0.209$	48	0.534	9.1–28.0
Gymnocanthus galeatus (armorhead sculpin)	BS	$WT = 0.0100\ SL^{3.196}$	29	0.939	$SL = 1.75\ (OL) + 0.82$ $SE_\beta = 0.360$	28	0.476	7.3–15.0
Hippoglossoides elassodon (flathead sole)	WA	$WT = 0.0078\ FL^{3.041}$	99	0.984	$FL = 4.63\ (OL) - 0.71$	40	0.947	17.2–30.5
Hyperprosopon argenteum (walleye surfperch)	OR, CA	$WT = 0.0116\ SL^{3.361}$	23	0.996	$SL = 2.57\ (OL) - 2.83$ $SE_\beta = 0.063$	24	0.987	5.6–20.7
Hypomesus pretiosus (surf smelt)	OR	$WT = 0.0044\ SL^{3.345}$	42	0.986	$SL = 3.61\ (OL) - 0.63$ $SE_\beta = 0.204$	25	0.932	6.9–15.4
Hypsopsetta guttulata (diamond turbot)	CA	$WT = 0.0853\ SL^{2.664}$	14	0.967	$SL = 4.89\ (OL) - 0.29$ $SE_\beta = 0.543$	18	0.835	7.1–26.9
Leptocottus armatus (Pacific staghorn sculpin)	WA,OR,CA	$WT = 0.0111\ SL^{3.229}$	51	0.990	$SL = 2.58\ (OL) - 2.26$ $SE_\beta = 0.092$	62	0.928	3.7–22.5
Lycodes brevipes (shortfin eelpout)	BS	$WT = 0.0195\ FL^{2.522}$	56	0.826	$FL = 3.47\ (OL) + 4.83$ $SE_\beta = 0.430$	62	0.520	18.5–27.6
Lycodes cortezianus (bigfin eelpout)	WA,CA	$WT = 0.0018\ SL^{3.245}$	40	0.993	$SL = 10.96\ (OL) - 21.82$	41	0.742	11.5–44.7
Lycodes palearis (wattled eelpout)	BS	$WT = 0.0007\ FL^{3.483}$	25	0.913	$FL = 5.22\ (OL) + 12.42$ $SE_\beta = 1.773$	24	0.283	32.0–47.0
Lycodopsis pacifica (blackbelly eelpout)	GA	$WT = 0.0018\ SL^{3.302}$	22	0.954	$SL = 3.82\ (OL) + 4.89$ $SE_\beta = 0.719$	20	0.610	14.4–23.3
Mallotus villosus (capelin)	BS	$WT = 0.0054\ SL^{3.160}$	39	0.717	$SL = 3.45\ (OL) + 3.62$ $SE_\beta = 0.417$	39	0.649	10.0–13.7
Merluccius productus (Pacific hake)	OR	$WT = 0.0081\ SL^{2.966}$	75	0.933	$SL = 2.04\ (OL) + 0.96$ $SE_\beta = 0.078$	86	0.891	26.3–54.4
Microgadus proximus (Pacific tomcod)	WA,OR	$WT = 0.0064\ SL^{3.191}$	80	0.988	$SL = 1.77\ (OL) - 3.51$ $SE_\beta = 0.048$	101	0.932	6.1–28.3
Microstomus pacificus (Dover sole)	OR	$WT = 0.0094\ SL^{3.092}$	101	0.854	$SL = 3.72\ (OL) + 6.97$ $SE_\beta = 0.301$	117	0.587	7.6–37.8
Oncorhynchus kisutch (coho salmon)	OR	$WT = 0.0103\ SL^{3.092}$	43	0.989	$SL = 16.31\ (OL) - 40.74$ $SE_\beta = 2.138$	46	0.569	12.5–58.4
Oncorhynchus mykiss (rainbow trout)	OR	$WT = 0.0275\ SL^{2.895}$	18	0.905	$SL = 16.28\ (OL) - 38.14$ $SE_\beta = 1.381$	39	0.790	12.0–26.1
Ophiodon elongatus (lingcod)	OR,CA	$WT = 0.0023\ SL^{3.567}$	10	0.620	$SL = 8.23\ (OL) - 8.20$ $SE_\beta = 0.315$	35	0.722	5.3–71.9
Osmerus mordax (rainbow smelt)	BS	$WT = 0.0038\ SL^{3.278}$	32	0.819	$SL = 2.69\ (OL) + 0.32$ $SE_\beta = 0.292$	29	0.759	13.0–17.9

Continued

Table 2 (continued)

Species (common name)	Location[1]	Fish weight/fish length			Fish length/otolith length			Range
		Equation	N	r^2	Equation	N	r^2	
Phanerodon furcatus (white seaperch)	OR,CA	$WT = 0.0213\ SL^{3.086}$	17	0.997	$SL = 2.33\ (OL) - 2.15$ $SE_\beta = 0.109$	46	0.912	4.5–23.5
Platichthys stellatus (starry flounder)	OR,CA	$WT = 0.0107\ SL^{3.268}$	25	0.985	$SL = 3.35\ (OL) + 0.23$ $SE_\beta = 0.303$	30	0.814	8.2–37.3
Pleurogrammus monopterygius (Atka mackerel)	BS	$WT = 0.0034\ FL^{3.401}$	16	0.987	$FL = 8.40\ (OL) - 4.99$	13	0.864	19.0–35.0
Pleuronectes asper (yellowfin sole)	BS	$WT = 0.0024\ SL^{3.605}$	7	0.859	$SL = 2.17\ (OL) + 10.65$	26	0.638	22.7–30.7
Pleuronectes bilineatus (rock sole)	BS	$WT = 0.0112\ FL^{2.997}$	83	0.931	$FL = 6.16\ (OL) - 6.97$	29	0.841	7.3–32.0
Pleuronectes vetulus (English sole)	WA,OR,CA	$WT = 0.0163\ SL^{2.939}$	98	0.995	$SL = 3.82\ (OL) - 2.76$ $SE_\beta = 0.059$	151	0.965	3.5–36.1
Podothecus acipenserinus (sturgeon poacher)	BS,WA, OR	$WT = 0.0030\ SL^{3.233}$	93	0.956	$SL = 6.58\ (OL) - 6.21$ $SE_\beta = 0.303$	92	0.840	7.3–25.7
Porichthys notatus (plainfin midshipman)	WA,OR, CA	$WT = 0.0207\ SL^{2.916}$	78	0.967	$SL = 2.80\ (OL) - 2.59$ $SE_\beta = 0.090$	80	0.926	4.0–26.6
Psettichthys melanostictus (sand sole)	WA,OR, CA	$WT = 0.0052\ SL^{3.441}$	25	0.983	$SL = 5.06\ (OL) - 3.18$ $SE_\beta = 0.256$	26	0.942	8.1–31.0
Rhacochilus vacca (pile perch)	WA,OR	$WT = 0.0182\ SL^{3.218}$	46	0.997	$SL = 3.35\ (OL) - 8.19$ $SE_\beta = 0.098$	45	0.965	6.2–33.5
Sebastes melanops (black rockfish)	OR	$WT = 0.1225\ SL^{2.499}$	21	0.585	$SL = 2.23\ (OL) - 1.48$ $SE_\beta = 0.181$	53	0.749	12.2–42.0
Sebastes paucispinis (bocaccio)	OR	No data available			$SL = 2.41\ (OL) + 0.14$ $SE_\beta = 0.398$	13	0.769	8.4–41.8
Sebastolobus alascanus (shortspine thornyhead)	OR	$WT = 0.0102\ SL^{3.239}$	69	0.988	$SL = 2.31\ (OL) - 3.71$ $SE_\beta = 0.150$	51	0.828	15.1–49.2
Sebastolobus altivelis (longspine thornyhead)	OR	$WT = 0.0155\ SL^{3.113}$	13	0.997	$SL = 4.94\ (OL) - 27.50$ $SE_\beta = 0.652$	13	0.839	15.8–55.5
Spirinchus thaleichthys (longfin smelt)	OR	$WT = 0.0288\ SL^{2.531}$	50	0.854	$SL = 2.64\ (OL) - 0.20$ $SE_\beta = 0.142$	50	0.878	7.3–11.6
Symphurus atricauda (California tonguefish)	CA	$WT = 0.0074\ SL^{3.136}$	32	0.789	$SL = 3.56\ (OL) + 4.64$ $SE_\beta = 0.563$	48	0.464	6.5–15.2
Thaleichthys pacificus (eulachon)	OR	$WT = 0.0077\ SL^{3.075}$	129	0.884	$SL = 4.71\ (OL) - 2.70$ $SE_\beta = 0.181$	102	0.871	10.5–19.8
Theragra chalcogramma (walleye pollock)	WA	$WT = 0.0043\ SL^{3.255}$	46	0.985	$SL = 2.24\ (OL) - 2.35$ $SE_\beta = 0.081$	44	0.948	12.6–43.2
Trachurus symmetricus (jack mackerel)	OR,CA	$WT = 0.0635\ SL^{2.556}$	18	0.761	Not significant	14	0.141	25.5–33.3
Trichodon trichodon (Pacific sandfish)	BS	$WT = 0.0170\ FL^{2.953}$	19	0.971	$FL = 6.06\ (OL) - 4.57$	17	0.684	15.0–25.0
Zalembius rosaceus (pink seaperch)	CA	$WT = 0.0199\ SL^{3.102}$	48	0.841	$SL = 1.88\ (OL) - 0.07$ $SE_\beta = 0.079$	56	0.912	7.0–12.8

[1] Location of capture: BS = Bering Sea, CA = California, GA = Gulf of Alaska, OR = Oregon, WA = Washington.

Brown and Mate, 1983; Harvey, 1987). Specific guides or keys to fish otoliths also have been published (Morrow, 1979; Harkonen, 1986; Hecht, 1987; Smale et al., 1995).

Generally, standard length of fishes is linearly related to otolith length. Predicting size of fishes (length and weight) can be accomplished with fair reliability on the basis of otolith length. This relationship, however, is not always reliable. Otolith length typically is linearly related to length of the fish until the fish reaches maximum size; thereafter, the otolith increases only in thickness (Blacker, 1974; Williams and Bedford, 1974). Otolith lengths of larval and juvenile fishes may increase in a curvilinear fashion relative to fish length for some species, such as sockeye salmon (*Oncorhynchus nerka*; West and Larkin, 1987) and walleye pollock (*Theragra chalcogramma*; Nishimura and Yamada, 1988). The relationship between otolith length and fish length may be dependent on the growth rate of the fish, as was reported for striped bass (*Morone saxatilis*; Secor and Dean, 1989). Additionally, the relationship between otolith length and fish length may be described by multiple linear lines with inflection points (e.g. gadids; Frost and Lowry, 1981). Multiple linear relationships may result from different growth stanzas (Laidig et al., 1991). These results indicate that size of fish should only be estimated over the size distribution sampled, and that all length intervals should be sampled properly with the appropriate statistical model.

Estimating size of consumed fishes from measurements of otoliths recovered in stomachs or feces may be biased because of partial or complete digestion of otoliths (Jobling and Breiby, 1986; Jobling, 1987). For instance, size of fish eaten by the harbor seal (*Phoca vitulina*) may be underestimated by 16–44% (Harvey, 1989). Similar results have been reported for many pinnipeds (Hawes, 1983; da Silva and Neilson, 1985; Murie and Lavigne, 1986). Although a rough estimate of these errors may be obtained from controlled experiments, the amount of digestion may be species-specific, requiring numerous tests to document all forms of bias. There also may be differences between the sexes in the relationship between fish size and otolith size, something we did not test. Researchers using otoliths to determine number and size of fish eaten, therefore, should realize the limitations of this technique.

Fish size-otolith size relationships will be useful for researchers examining food habits of piscivores and size of fish in archaeological samples. Many more species and sizes of fish should be sampled to cover the full range of fishes involved in these studies.

Acknowledgments

We thank students and coworkers at Oregon State University, Moss Landing Marine Laboratories, and the National Marine Mammal Laboratory for their assistance in removing and measuring fish otoliths. In particular, Mary Yoklavich helped enter data, checked data for accuracy, and reviewed the manuscript. Daniel Kimura, Mark Lowry, Beth Sinclair, and Jay Orr also provided thoughtful comments on the manuscript. This work was funded, in part, by the National Marine Mammal Laboratory, National Marine Fisheries Service, Seattle, Washington.

Literature Cited

Ainley, D. G., D. W. Anderson, and P. R. Kelly.
1981. Feeding ecology of marine cormorants in southwestern North America. Condor 83:120–131.

Antonelis, G. A., Jr., C. H. Fiscus, and R. L. DeLong.
1984. Spring and summer prey of California sea lions, *Zalophus californianus*, at San Miguel Island, California, 1978–79. Fish. Bull. 82:67–76.

Bailey, K. M., and D. G. Ainley.
1982. The dynamics of California sea lion predation on Pacific hake. Fish. Res. 1:163–176.

Blacker, R. W.
1974. Recent advances in otolith studies. *In* F. R. Harden Jones (ed.), Sea fisheries research, p. 67–90. John Wiley and Sons, N.Y.

Boehlert, G. W.
1985. Using objective criteria and multiple regression models for age determination in fishes. Fish. Bull. 83:103–117.

Boehlert, G. W., and M. M. Yoklavich.
1984. Variability in age estimates in *Sebastes* as a function of methodology, different readers, and different laboratories. Calif. Fish Game 70:210–224.

Brown, R. B., and B. R. Mate.
1983. Abundance, movements, and feeding habits of harbor seals, *Phoca vitulina*, at Netarts and Tillamook Bays, Oregon. Fish. Bull. 81:291–301.

Casteel, R.
1974. Use of Pacific salmon otoliths for estimating fish size, with a note on the size of late Pleistocene and Pliocene salmonids. Northwest Sci. 48:175–179.
1976. Fish remains in archaeology and paleo-environmental studies. Acad. Press, N.Y., 180 p.

Chilton, D. E., and R. J. Beamish.
1982. Age determination methods for fishes studied by the groundfish program at the Pacific Biological Station. Fish. Aquat. Sci. Spec. Publ. 60, 102 p.

da Silva, J., and J. D. Neilson.
1985. Limitations of using otoliths recovered in scats to estimate prey consumption in seals. Can. J. Fish. Aquat. Sci. 42:1439–1442.

Fitch, J. E.
1964. The fish fauna of the Playa del Rey locality, a southern California marine Pleistocene deposit. Los Angel. Cty. Mus. Contrib. Sci. 82:3–35.
1968. Otoliths and other fish remains from the Timms Point silt (early Pleistocene) at San Pedro, California. Los Angel. Cty. Mus. Contrib. Sci. 146:1–29.
1969. Fossil records of certain schooling fishes of the California current system. CalCOFI Rep. 13:71–80.

Fitch, J. E., and R. L. Brownell Jr.
1968. Fish otoliths in cetacean stomachs and their importance in interpreting feeding habits. J. Fish. Res. Board Can. 25:2561–2574.

Frost, K. J., and L. F. Lowry.
1981. Trophic importance of some marine gadids in northern Alaska and their body-otolith size relationships. Fish. Bull. 79: 187–192.

Gamboa, D. A.
1991. Otolith size versus weight and body-length relationships for eleven fish species of Baja California, Mexico. Fish Bull. 89:701–706.

Harkonen, T.
1986. Guide to the otoliths of the bony fishes of the northeast Atlantic. Danbiu Aps, Denmark, 256 p.

Harvey, J. T.
1987. Population dynamics, annual food consumption, movements, and dive behaviors of harbor seals, *Phoca vitulina*, in Oregon. Ph.D. dissert., Oregon State Univ., Corvallis, 177 p.
1989. Assessment of errors associated with harbour seal (*Phoca vitulina*) faecal sampling. J. Zool. (Lond.) 219:101–111.

Hawes, S. D.
1983. An evaluation of California sea lion scat samples as indicators of prey importance. M.S. thesis, San Francisco State Univ., CA, 50 p.

Hecht, T.
1977. Otoliths: what they are and their scientific importance. Eastern Cape Naturalist 61:18–20.
1987. A guide to the otoliths of southern ocean fishes. J. Antarctic Res. 17:1–87.

Jobling, M.
1987. Marine mammal faeces as indicators of prey importance—a source of error in bioenergetics studies. Sarsia 72: 255–260.

Jobling, M., and A. Breiby.
1986. The use and abuse of fish otoliths in studies of feeding habits of marine piscivores. Sarsia 71:265–274.

Laidig, T. E., S. Ralston, and J. R. Bence.
1991. Dynamics of growth in the early life history of shortbelly rockfish *Sebastes jordani*. Fish. Bull. 89:611–621.

McKern, J. L., H. F. Horton, and K. V. Koski.
1974. Development of steelhead trout (*Salmo gairdneri*) otoliths, and their use for age analysis and for separating summer from winter races and wild from hatchery stocks. J. Fish. Res. Board Can. 31:1420–1426.

Messieh, S. N.
1972. Use of otoliths in identifying herring stocks in the southern Gulf of St. Lawrence and adjacent waters. J. Fish. Res. Board Can. 29:1113–1118.

Morrow, J. E.
1979. Preliminary keys to otoliths of some adult fishes of the Gulf of Alaska, Bering Sea, and Beaufort Sea. U.S. Dep. Commer., NOAA Tech. Rep. NMFS Circ. 420, 32 p.

Murie, D. J., and D. M. Lavigne.
1986. Interpretation of otoliths in stomach content analyses of phocid seals: quantifying fish consumption. Can. J. Zool. 64:1152–1157.

Nishimura, A., and J. Yamada.
1988. Geographical differences in early growth of walleye pollock *Theragra chalcogramma*, estimated by back-calculation of otolith daily growth increments. Mar. Biol. 97:459–465.

Pitcher, K. W.
1980. Stomach contents and feces as indicators of harbor seal, *Phoca vitulina*, foods in the Gulf of Alaska. Fish. Bull. 78: 797–798.

Ricker, W. E.
1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191, 382 p.

Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. Lea, and W. B. Scott.
1991. Common and scientific names of fishes from the United States and Canada. Am. Fish. Soc. Spec. Publ. 20, 183 p.

Saila, S. B., C. W. Recksiek, and M. H. Prager.
1988. Basic fishery science programs. Elsevier, Amsterdam, 230 p.

Secor, D. H., and J. M. Dean.
1989. Somatic growth effects on the otolith-fish size relationship in young pond-reared striped bass, *Morone saxatilis*. Can. J. Fish. Aquat. Sci. 46:113–121.

Smale, M. J., G. Watson, and T. Hecht.
1995. Otolith atlas of southern African marine fishes. Ichthyol. Monogr. J.L.B. Smith Inst. Ichthyology 1:1–253.

Southward, G. M.
1962. A method of calculating body lengths from otolith measurements for Pacific halibut and its application to portlock-albatross grounds data between 1935 and 1957. J. Fish. Res. Board Can. 19:339–362.

Southward, G. M., and D. G. Chapman.
1965. Utilization of Pacific halibut stocks: study of Bertlanffy's growth equation. Rep. Int. Pac. Halibut Comm. 39:1–33.

Spratt, J. D.
1975. Growth rate of the northern anchovy, *Engraulis mordax*, in southern California waters, calculated from otoliths. Calif. Fish Game 61:116–126.

Summerfelt, R. C., and G. E. Hall (eds.).
1987. The age and growth of fish. Iowa State Univ. Press, Ames, 544 p.

Templemann, W., and H. J. Squires.
1956. Relationship of otolith lengths and weights in the haddock, *Melanogrammus aeglefinus* (L.), to the growth of the fish. J. Fish. Res. Board Can. 13:467–487.

Treacy, S. D., and T. W. Crawford.
1981. Retrieval of otoliths and statoliths from gastrointestinal contents and scats of marine mammals. J. Wildl. Manage. 45:990–993.

Trippel, E. A., and F. W. H. Beamish.
1987. Characterizing piscivory from ingested remains. Trans. Am. Fish. Soc. 116:773–776.

Trout, G. C.
1954. Otolith growth of the Barents Sea cod. Rapp. P-v. Reun. Cons. Int. Explor. Mer 150:297–299.

West, C. J., and P. A. Larkin.
1987. Evidence of size-selective mortality of juvenile sockeye salmon (*Oncorhynchus nerka*) in Babine Lake, British Columbia. Can. J. Fish. Aquat. Sci. 44:712–721.

Williams, T., and B. C. Bedford.
1974. The use of otoliths for age determination. *In* T. B. Bagenal (ed.), The ageing of fish, p. 114–123. Unwin Brothers, Surrey, England.

Wyllie Echeverria, T.
1987. Relationship of otolith length to total length in rockfishes from northern and central California. Fish. Bull. 85:383–387.

Figure 1

Otolith images of 76 species of fishes from the eastern North Pacific, listed in taxonomic order (Robins et al., 1991). For species with extreme variability in otolith morphology, multiple images are provided. For each species, the scientific and common names, position and size of the otolith pictured, and length and mass of the fish from which the otolith was removed are given. The regression relationships between (1) weight (WT in grams) and fish length (SL in cm) and (2) fish length (SL or FL in cm) and otolith length (OL in mm) are provided for each species. The coefficient of determination is r^2.

CLUPEIDAE



Alosa sapidissima
(American shad)

Left otolith; length (mm): 3.6
Fish length (cm): 27.3
Fish weight (g): 320.5

Regression equations:

Length: SL = 11.46 (OL) – 11.08 $r^2 = 0.960$
Weight: WT = 0.0135 $SL^{3.046}$ $r^2 = 0.997$



Clupea pallasi
(Pacific herring)

Right otolith; length (mm): 3.3
Fish length (cm): 17.7
Fish weight (g): 71.6

Regression equations:

Length: SL = 5.24 (OL) – 1.85 $r^2 = 0.934$
Weight: WT = 0.0044 $SL^{3.398}$ $r^2 = 0.976$

ENGRAULIDAE



Engraulis mordax
(northern anchovy)

Right otolith; length (mm): 3.5
Fish length (cm): 8.0
Fish weight (g): 7.8

Regression equations:

Length: SL = 2.28 (OL) + 0.85 $r^2 = 0.694$
Weight: WT = 0.0485 $SL^{2.413}$ $r^2 = 0.807$



Engraulis mordax
(northern anchovy)

Left otolith; length (mm): 3.4
Fish length (cm): 11.9
Fish weight (g): 12.8

Regression equations:

Length: SL = 2.28 (OL) + 0.85 $r^2 = 0.694$
Weight: WT = 0.0485 $SL^{2.413}$ $r^2 = 0.807$

OSMERIDAE



Allosmerus elongatus
(whitebait smelt)

Right otolith; length (mm): 3.0
Fish length (cm): 9.3
Fish weight (g): 5.9

Regression equations:

Length: SL = 2.11 (OL) + 3.02 $r^2 = 0.838$
Weight: WT = 0.0063 $SL^{3.233}$ $r^2 = 0.893$

Hypomesus pretiosus
(surf smelt)

Left otolith; length (mm): 3.8
Fish length (cm): 12.6
Fish weight (g): 20.1

Regression equations:

Length: SL = 3.61 (OL) – 0.63 $r^2 = 0.932$
Weight: WT = 0.0044 $SL^{3.345}$ $r^2 = 0.986$



Mallotus villosus
(capelin)

Right otolith; length (mm): 2.5
Fish length (cm): 11.6
Fish weight (g): 12.0

Regression equations:

Length: SL = 3.45 (OL) + 3.62 $r^2 = 0.649$
Weight: WT = 0.0054 $SL^{3.160}$ $r^2 = 0.717$



Osmerus mordax
(rainbow smelt)

Left otolith; length (mm): 5.1
Fish length (cm): 13.0
Fish weight (g): 17.9

Regression equations:

Length: SL = 2.69 (OL) + 0.32 $r^2 = 0.759$
Weight: WT = 0.0038 $SL^{3.278}$ $r^2 = 0.819$

OSMERIDAE (cont.)



Spirinchus thaleichthys
(longfin smelt)

Left otolith; length (mm): 3.8
Fish length (cm): 9.9
Fish weight (g): 8.8

Regression equations:

Length: SL = 2.64 (OL) – 0.20 $r^2 = 0.878$
Weight: WT = 0.0288 $SL^{2.531}$ $r^2 = 0.854$



Thaleichthys pacificus
(eulachon)

Left otolith; length (mm): 4.1
Fish length (cm): 16.5
Fish weight (g): 32.7

Regression equations:

Length: SL = 4.71 (OL) – 2.70 $r^2 = 0.871$
Weight: WT = 0.0077 $SL^{3.075}$ $r^2 = 0.884$

SALMONIDAE



Oncorhynchus kisutch
(coho salmon)

Right otolith; length (mm): 3.7
Fish length (cm): 18.0
Fish weight (g): 98.9

Regression equations:

Length: SL = 16.31 (OL) – 40.74 $r^2 = 0.569$
Weight: WT = 0.0103 $SL^{3.092}$ $r^2 = 0.989$

Oncorhynchus kisutch
(coho salmon)

Right otolith; length (mm): 3.3
Fish length (cm): 17.7
Fish weight (g): 66.6

Regression equations:

Length: SL = 16.31 (OL) – 40.74 $r^2 = 0.569$
Weight: WT = 0.0103 $SL^{3.092}$ $r^2 = 0.989$

SALMONIDAE (cont.)

Oncorhynchus mykiss
(rainbow trout)

Left otolith; length (mm): 5.0
Fish length (cm): 24.3
Fish weight (g): 320.9

Regression equations:

Length: SL = 16.28 (OL) – 38.14 $r^2 = 0.790$
Weight: WT = 0.0275 $SL^{2.895}$ $r^2 = 0.905$



Oncorhynchus mykiss
(rainbow trout)

Right otolith; length (mm): 4.8
Fish length (cm): 25.3
Fish weight (g): 315.6

Regression equations:

Length: SL = 16.28 (OL) – 38.14 $r^2 = 0.790$
Weight: WT = 0.0275 $SL^{2.895}$ $r^2 = 0.905$

Oncorhynchus nerka
(sockeye salmon)

Left otolith; length (mm): 3.0
Fish length (cm): 20
Fish weight (g): N/A

Regression equations:

Length: No data available
Weight: No data available



Oncorhynchus tshawytscha
(chinook salmon)

Left otolith; length (mm): 2.4
Fish length (cm): 7.6
Fish weight (g): N/A

Regression equations:

Length: No data available
Weight: No data available

GADIDAE

Eleginus gracilis
(Saffron cod; ventral view)

Left otolith; length (mm): 6.8
Fish length (cm): 9.2
Fish weight (g): 6.2

Regression equations:
Length: SL = 1.89 (OL) – 2.76 $r^2 = 0.960$
Weight: WT = 0.0039 $SL^{3.292}$ $r^2 = 0.990$



Eleginus gracilis
(saffron cod)

Left otolith; length (mm): 6.8
Fish length (cm): 9.2
Fish weight (g): 6.2

Regression equations:
Length: SL = 1.89 (OL) – 2.76 $r^2 = 0.960$
Weight: WT = 0.0039 $SL^{3.292}$ $r^2 = 0.990$



Gadus macrocephalus
(Pacific cod)

Right otolith; length (mm): 12.0
Fish length (cm): 29.5
Fish weight (g): 373.8

Regression equations:
Length: FL = 4.51 (OL) – 22.97 $r^2 = 0.883$
Weight: No data available

Merluccius productus
(Pacific hake)

Right otolith; length (mm): 22.0
Fish length (cm): 42.0
Fish weight (g): 659.0

Regression equations:
Length: SL = 2.04 (OL) + 0.96 $r^2 = 0.891$
Weight: WT = 0.0081 $SL^{2.966}$ $r^2 = 0.933$

GADIDAE (cont.)



Microgadus proximus
(Pacific tomcod)

Right otolith; length (mm): 12.3
Fish length (cm): 18.2
Fish weight (g): 59.8

Regression equations:
Length: SL = 1.77 (OL) – 3.51 $r^2 = 0.932$
Weight: WT = 0.0064 $SL^{3.191}$ $r^2 = 0.988$



Theragra chalcogramma
(walleye pollock)

Left otolith; length (mm): 16.0
Fish length (cm): 33.5
Fish weight (g): 394.0

Regression equations:
Length: SL = 2.24 (OL) – 2.35 $r^2 = 0.948$
Weight: WT = 0.0043 $SL^{3.255}$ $r^2 = 0.985$

MACROURIDAE



Coryphaenoides acrolepis
(Pacific rattail)

Right otolith; length (mm): 10.0
Fish length (cm): 36.0
Fish weight (g): 148.8

Regression equations:
Length: SL = 3.44 (OL) – 3.23 $r^2 = 0.926$
Weight: WT = 0.0016 $SL^{3.209}$ $r^2 = 0.921$

OPHIDIIDAE



Chilara taylori
(spotted cusk-eel)

Left otolith; length (mm): 7.6
Fish length (cm): 21.8
Fish weight (g): 46.6

Regression equations:
Length: SL = 2.51 (OL) + 2.15 $r^2 = 0.730$
Weight: WT = 0.0004 $SL^{3.761}$ $r^2 = 0.964$

BATRACHOIDIDAE



Porichthys notatus
(plainfin midshipman)

Left otolith; length (mm): 9.1
Fish length (cm): 22.5
Fish weight (g): 215.1

Regression equations:

Length: SL = 2.80 (OL) – 2.59 $r^2 = 0.926$
Weight: WT = 0.0207 $SL^{2.916}$ $r^2 = 0.967$

ATHERINIDAE

Atherinops affinis
(topsmelt)

Right otolith; length (mm): 2.4
Fish length (cm): 9.9
Fish weight (g): 10.2

Regression equations:

Length: SL = 3.72 (OL) + 0.55 $r^2 = 0.891$
Weight: WT = 0.1698 $SL^{1.733}$ $r^2 = 0.429$



Atherinops affinis
(topsmelt)

Left otolith; length (mm): 3.2
Fish length (cm): 11.8
Fish weight (g): 13.2

Regression equations:

Length: SL = 3.72 (OL) + 0.55 $r^2 = 0.891$
Weight: WT = 0.1698 $SL^{1.733}$ $r^2 = 0.429$



Atherinopsis californiensis
(jacksmelt)

Right otolith; length (mm): 6.1
Fish length (cm): 28.7
Fish weight (g): 260.0

Regression equations:

Length: SL = 4.85 (OL) – 2.46 $r^2 = 0.950$
Weight: WT = 0.0049 $SL^{3.228}$ $r^2 = 0.968$

SCORPAENIDAE



Sebastes auriculatus
(brown rockfish)

Left otolith; length (mm): 3.2
Fish length (cm): 6.7
Fish weight (g): 6.9

Regression equations:
Length: SL = 3.32 (OL) – 5.30 $r^2 = 0.940$
Weight: No data available



Sebastes constellatus
(starry rockfish)

Right otolith; length (mm): 12.3
Fish length (cm): N/A
Fish weight (g): N/A

Regression equations:
Length: SL = 2.53 (OL) – 3.75 $r^2 = 0.960$
Weight: No data available



Sebastes crameri
(darkblotched rockfish)

Left otolith; length (mm): 13.9
Fish length (cm): 31.0
Fish weight (g): 1121.5

Regression equations:
Length: SL = 2.81 (OL) – 2.71 $r^2 = 0.940$
Weight: No data available



Sebastes diploproa
(splitnose rockfish)

Right otolith; length (mm): 15.1
Fish length (cm): N/A
Fish weight (g): N/A

Regression equations:
Length: SL = 2.26 (OL) – 1.29 $r^2 = 0.960$
Weight: No data available

SCORPAENIDAE (cont.)



Sebastes flavidus
(yellowtail rockfish)

Right otolith; length (mm): 12.2
Fish length (cm): 24.5
Fish weight (g): N/A

Regression equations:
Length: SL = 2.65 (OL) – 1.09 $r^2 = 0.880$
Weight: No data available



Sebastes maliger
(quillback rockfish)

Left otolith; length (mm): 13.0
Fish length (cm): 32.5
Fish weight (g): N/A

Regression equations:
Length: SL = 2.99 (OL) – 5.31 $r^2 = 0.860$
Weight: No data available

Sebastes melanops
(black rockfish)

Right otolith; length (mm): 12.4
Fish length (cm): 29.0
Fish weight (g): N/A

Regression equations:
Length: SL = 2.23 (OL) – 1.48 $r^2 = 0.749$
Weight: WT = 0.1225 $SL^{2.499}$ $r^2 = 0.585$



Sebastes miniatus
(vermillion rockfish)

Right otolith; length (mm): 15.1
Fish length (cm): 35.5
Fish weight (g): N/A

Regression equations:
Length: SL = 2.94 (OL) – 5.67 $r^2 = 0.930$
Weight: No data available

SCORPAENIDAE (cont.)



Sebastes mystinus
(blue rockfish)

Right otolith; length (mm): 12.8
Fish length (cm): 29.0
Fish weight (g): N/A

Regression equations:
Length: SL = 2.98 (OL) – 1.82 $r^2 = 0.830$
Weight: No data available



Sebastes nebulosus
(China rockfish)

Right otolith; length (mm): 10.7
Fish length (cm): N/A
Fish weight (g): N/A

Regression equations:
Length: SL = 2.52 (OL) + 3.30 $r^2 = 0.790$
Weight: No data available



Sebastes paucispinis
(boccacio)

Left otolith; length (mm): 14.0
Fish length (cm): 39.8
Fish weight (g): 1191.8

Regression equations:
Length: SL = 2.41 (OL) + 0.14 $r^2 = 0.769$
Weight: No data available



Sebastes pinniger
(canary rockfish)

Right otolith; length (mm): 16.3
Fish length (cm): 31.8
Fish weight (g): N/A

Regression equations:
Length: SL = 2.94 (OL) – 8.51 $r^2 = 0.920$
Weight: No data available

SCORPAENIDAE (cont.)



Sebastes ruberrimus
(yellow eye rockfish)

Left otolith; length (mm): 14.4
Fish length (cm): 34.0
Fish weight (g): N/A

Regression equations:
Length: SL = 3.13 (OL) – 7.62 $r^2 = 0.920$
Weight: No data available



Sebastolobus alascanus
(shortspine thornyhead)

Right otolith; length (mm): 14.1
Fish length (cm): 27.7
Fish weight (g): 515.6

Regression equations:
Length: SL = 2.31 (OL) – 3.71 $r^2 = 0.828$
Weight: WT = 0.0102 $SL^{3.239}$ $r^2 = 0.988$



Sebastolobus altivelis
(longspine thornyhead)

Right otolith; length (mm): 10.9
Fish length (cm): 24.5
Fish weight (g): 328.8

Regression equations:
Length: SL = 4.94 (OL) – 27.50 $r^2 = 0.839$
Weight: WT = 0.0155 $SL^{3.113}$ $r^2 = 0.997$

ANOPLOPOMATIDAE

Anoplopoma fimbria
(sablefish)

Left otolith; length (mm): 6.2
Fish length (cm): 36.4
Fish weight (g): 541.7

Regression equations:

Length: SL = 5.28 (OL) + 1.62 $r^2 = 0.955$
Weight: WT = 0.0163 SL$^{2.902}$ $r^2 = 0.993$



Anoplopoma fimbria
(sablefish)

Right otolith; length (mm): 7.7
Fish length (cm): 41.6
Fish weight (g): 868.1

Regression equations:

Length: SL = 5.28 (OL) + 1.62 $r^2 = 0.955$
Weight: WT = 0.0163 SL$^{2.902}$ $r^2 = 0.993$

HEXAGRAMMIDAE



Ophiodon elongatus
(lingcod)

Right otolith; length (mm): 9.1
Fish length (cm): 62.0
Fish weight (g): N/A

Regression equations:

Length: SL = 8.23 (OL) – 8.20 $r^2 = 0.722$
Weight: WT = 0.0023 SL$^{3.567}$ $r^2 = 0.620$



Ophiodon elongatus
(lingcod)

Left otolith; length (mm): 5.3
Fish length (cm): 38.2
Fish weight (g): N/A

Regression equations:

Length: SL = 8.23 (OL) – 8.20 $r^2 = 0.722$
Weight: WT = 0.0023 SL$^{3.567}$ $r^2 = 0.620$

HEXAGRAMMIDAE (cont.)



Ophiodon elongatus
(lingcod)

Left otolith; length (mm): 9.0
Fish length (cm): 62.0
Fish weight (g): N/A

Regression equations:

Length: SL = 8.23 (OL) – 8.20 $r^2 = 0.722$
Weight: WT = 0.0023 SL$^{3.567}$ $r^2 = 0.620$



Pleurogrammus monopterygius
(Atka mackerel)

Right otolith; length (mm): 3.5
Fish length (cm): 21.2
Fish weight (g): 108.8

Regression equations:

Length: FL = 8.40 (OL) – 4.99 $r^2 = 0.864$
Weight: WT = 0.0034 FL$^{3.401}$ $r^2 = 0.987$

COTTIDAE



Chitonotus pugetensis
(roughback sculpin)

Right otolith; length (mm): 2.9
Fish length (cm): 6.0
Fish weight (g): 3.3

Regression equations:

Length: SL = 3.37 (OL) – 4.52 $r^2 = 0.857$
Weight: WT = 0.0217 SL$^{2.871}$ $r^2 = 0.960$

Dasycottus setiger
(spinyhead sculpin)

Left otolith; length (mm): 8.5
Fish length (cm): N/A
Fish weight (g): N/A

Regression equations:

Length: FL = 3.11 (OL) – 7.03 $r^2 = 0.655$
Weight: No data available

COTTIDAE (cont.)



Dasycottus setiger
(spinyhead sculpin)

Right otolith; length (mm): 8.3
Fish length (cm): N/A
Fish weight (g): N/A

Regression equations:
Length: FL = 3.11 (OL) – 7.03 $r^2 = 0.655$
Weight: No data available



Gymnocanthus galeatus
(armorhead sculpin)

Right otolith; length (mm): 6.6
Fish length (cm): 13.4
Fish weight (g): 43.5

Regression equations:
Length: SL = 1.75 (OL) + 0.82 $r^2 = 0.476$
Weight: WT = 0.0100 $SL^{3.196}$ $r^2 = 0.939$



Gymnocanthus galeatus
(armorhead sculpin)

Right otolith; length (mm): 6.5
Fish length (cm): 11.3
Fish weight (g): 22.4

Regression equations:
Length: SL = 1.75 (OL) + 0.82 $r^2 = 0.476$
Weight: WT = 0.0100 $SL^{3.196}$ $r^2 = 0.939$



Leptocottus armatus
(Pacific staghorn sculpin)

Right otolith; length (mm): 5.2
Fish length (cm): 11.5
Fish weight (g): 30.5

Regression equations:
Length: SL = 2.58 (OL) – 2.26 $r^2 = 0.928$
Weight: WT = 0.0111 $SL^{3.229}$ $r^2 = 0.990$

COTTIDAE (cont.)



Leptocottus armatus
(Pacific staghorn sculpin)

Left otolith; length (mm): 5.8
Fish length (cm): 13.6
Fish weight (g): 46.5

Regression equations:

Length: SL = 2.58 (OL) – 2.26 $r^2 = 0.928$
Weight: WT = 0.0111 SL$^{3.229}$ $r^2 = 0.990$



Leptocottus armatus
(Pacific staghorn sculpin)

Left otolith; length (mm): 6.1
Fish length (cm): 13.5
Fish weight (g): 47.8

Regression equations:

Length: SL = 2.58 (OL) – 2.26 $r^2 = 0.928$
Weight: WT = 0.0111 SL$^{3.229}$ $r^2 = 0.990$

AGONIDAE



Podothecus acipenserinus
(sturgeon poacher)

Right otolith; length (mm): 4.2
Fish length (cm): 19.1
Fish weight (g): 54.0

Regression equations:

Length: SL = 6.58 (OL) – 6.21 $r^2 = 0.840$
Weight: WT = 0.0030 SL$^{3.233}$ $r^2 = 0.956$

CARANGIDAE



Trachurus symmetricus
(Jack mackerel)

Right otolith; length (mm): 7.6
Fish length (cm): 26.8
Fish weight (g): 266.0

Regression equations:

Length: Not significant $r^2 = 0.141$
Weight: WT = 0.0635 SL$^{2.556}$ $r^2 = 0.761$

SCIAENIDAE



Genyonemus lineatus **(white croaker)**

Left otolith; length (mm): 10.2
Fish length (cm): 20.5
Fish weight (g): 171.8

Regression equations:

Length: SL = 1.52 (OL) + 4.66 $r^2 = 0.534$
Weight: WT = 0.0550 $SL^{2.700}$ $r^2 = 0.767$

Genyonemus lineatus **(lateral view) (white croaker)**

Left otolith; length (mm): 10.2
Fish length (cm): 20.5
Fish weight (g): 171.8

Regression equations:

Length: SL = 1.52 (OL) + 4.66 $r^2 = 0.534$
Weight: WT = 0.0550 $SL^{2.700}$ $r^2 = 0.767$

EMBIOTOCIDAE



Cymatogaster aggregata **(shiner perch)**

Right otolith; length (mm): 6.3
Fish length (cm): 10.1
Fish weight (g): 35.7

Regression equations:

Length: SL = 1.74 (OL) – 0.52 $r^2 = 0.925$
Weight: WT = 0.0100 $SL^{3.515}$ $r^2 = 0.979$



Cymatogaster aggregata **(shiner perch)**

Left otolith; length (mm): 6.3
Fish length (cm): 10.7
Fish weight (g): 38.6

Regression equations:

Length: SL = 1.74 (OL) – 0.52 $r^2 = 0.925$
Weight: WT = 0.0100 $SL^{3.515}$ $r^2 = 0.979$

EMBIOTOCIDAE (cont.)



Embiotoca jacksoni
(black perch)

Right otolith; length (mm): 7.4
Fish length (cm): 15.6
Fish weight (g): 169.0

Regression equations:

Length: SL = 2.45 (OL) – 2.61 $r^2 = 0.947$
Weight: WT = 0.0282 SL$^{3.148}$ $r^2 = 0.992$

Embiotoca lateralis
(striped seaperch)

Right otolith; length (mm): 10.5
Fish length (cm): 24.2
Fish weight (g): 540.2

Regression equations:

Length: SL = 2.90 (OL) – 5.68 $r^2 = 0.990$
Weight: WT = 0.0329 SL$^{3.043}$ $r^2 = 0.998$



Hyperprosopon argenteum
(walleye surfperch)

Right otolith; length (mm): 7.9
Fish length (cm): 18.7
Fish weight (g): 213.0

Regression equations:

Length: SL = 2.57 (OL) – 2.83 $r^2 = 0.987$
Weight: WT = 0.0116 SL$^{3.361}$ $r^2 = 0.996$



Phanerodon furcatus
(white seaperch)

Right otolith; length (mm): 8.7
Fish length (cm): 23.5
Fish weight (g): 402.2

Regression equations:

Length: SL = 2.33 (OL) – 2.15 $r^2 = 0.912$
Weight: WT = 0.0213 SL$^{3.086}$ $r^2 = 0.997$

EMBIOTOCIDAE (cont.)



Rhacochilus vacca
(pile perch)

Right otolith; length (mm): 11.8
Fish length (cm): 31.2
Fish weight (g): 1326.40

Regression equations:

Length: SL = 3.35 (OL) – 8.19 $r^2 = 0.965$
Weight: WT = 0.0182 SL$^{3.218}$ $r^2 = 0.997$



Zalembius rosaceus
(pink seaperch)

Left otolith; length (mm): 4.7
Fish length (cm): 8.9
Fish weight (g): 19.0

Regression equations:

Length: SL = 1.88 (OL) – 0.07 $r^2 = 0.912$
Weight: WT = 0.0199 SL$^{3.102}$ $r^2 = 0.841$

BATHYMASTERIDAE



Bathymaster signatus
(searcher)

Left otolith; length (mm): 6.3
Fish length (cm): 21.0
Fish weight (g): 150.3

Regression equations:

Length: SL = 3.48 (OL) + 1.90 $r^2 = 0.883$
Weight: WT = 0.0038 SL$^{3.256}$ $r^2 = 0.991$

Bathymaster signatus
(searcher)

Left otolith; length (mm): 6.9
Fish length (cm): 23.3
Fish weight (g): 212.9

Regression equations:

Length: SL = 3.48 (OL) + 1.90 $r^2 = 0.883$
Weight: WT = 0.0038 SL$^{3.256}$ $r^2 = 0.991$

ZOARCIDAE

Lycodes brevipes
(shortfin eelpout)

Left otolith; length (mm): 5.1
Fish length (cm): 24.0
Fish weight (g): 61.3

Regression equations:

Length: FL = 3.47 (OL) + 4.83 $r^2 = 0.520$
Weight: WT = 0.0195 FL$^{2.522}$ $r^2 = 0.826$



Lycodes cortezianus
(bigfin eelpout)

Left otolith; length (mm): 4.9
Fish length (cm): 38.5
Fish weight (g): 286.8

Regression equations:

Length: SL = 10.96 (OL) – 21.82 $r^2 = 0.742$
Weight: WT = 0.0018 SL$^{3.245}$ $r^2 = 0.993$



Lycodopsis pacifica
(blackbelly eelpout)

Right otolith; length (mm): 3.6
Fish length (cm): 19.5
Fish weight (g): 30.6

Regression equations:

Length: SL = 3.82 (OL) + 4.89 $r^2 = 0.610$
Weight: WT = 0.0018 SL$^{3.302}$ $r^2 = 0.954$

TRICHODONTIDAE



Trichodon trichodon
(Pacific sandfish)

Right otolith; length (mm): 4.0
Fish length (cm): N/A
Fish weight (g): N/A

Regression equations:

Length: FL = 6.06 (OL) – 4.57 $r^2 = 0.684$
Weight: WT = 0.0170 FL$^{2.953}$ $r^2 = 0.971$

AMMODYTIDAE



Ammodytes hexapterus
(Pacific sand lance)

Right otolith; length (mm): 2.4
Fish length (cm): 11.7
Fish weight (g): 6.4

Regression equations:

Length: SL = 4.06 (OL) + 2.01 $r^2 = 0.433$
Weight: WT = 0.0063 SL$^{2.790}$ $r^2 = 0.913$

BOTHIDAE



Citharichthys sordidus
(Pacific sanddab)

Left otolith; length (mm): 6.3
Fish length (cm): 23.0
Fish weight (g): 175.8

Regression equations:

Length: SL = 2.87 (OL) + 3.29 $r^2 = 0.727$
Weight: WT = 0.0352 SL$^{2.710}$ $r^2 = 0.851$



Citharichthys sordidus
(Pacific sanddab)

Right otolith; length (mm): 5.8
Fish length (cm): 20.6
Fish weight (g): 119.5

Regression equations:

Length: SL = 2.87 (OL) + 3.29 $r^2 = 0.727$
Weight: WT = 0.0352 SL$^{2.710}$ $r^2 = 0.851$

PLEURONECTIDAE



Atheresthes stomias
(arrowtooth flounder)

Right otolith; length (mm): 9.0
Fish length (cm): 37.5
Fish weight (g): 662.6

Regression equations:

Length: FL = 4.75 (OL) – 2.96 $r^2 = 0.925$
Weight: WT = 0.0093 FL$^{2.999}$ $r^2 = 0.961$



Eopsetta exilis
(slender sole)

Left otolith; length (mm): 4.6
Fish length (cm): 19.6
Fish weight (g): 95.4

Regression equations:

Length: SL = 3.37 (OL) + 1.08 $r^2 = 0.771$
Weight: WT = 0.0058 SL$^{3.293}$ $r^2 = 0.974$



Eopsetta exilis
(slender sole)

Right otolith; length (mm): 4.6
Fish length (cm): 18.5
Fish weight (g): 61.0

Regression equations:

Length: SL = 3.37 (OL) + 1.08 $r^2 = 0.771$
Weight: WT = 0.0058 SL$^{3.293}$ $r^2 = 0.974$

Eopsetta jordani
(petrale sole)

Right otolith; length (mm): 7.3
Fish length (cm): 29.0
Fish weight (g): 502.7

Regression equations:

Length: SL = 4.85 (OL) – 4.81 $r^2 = 0.857$
Weight: WT = 0.0086 SL$^{3.231}$ $r^2 = 0.986$

PLEURONECTIDAE (cont.)



Errex zachirus
(rex sole)

Right otolith; length (mm): 6.6
Fish length (cm): 27.2
Fish weight (g): 171.0

Regression equations:

Length: SL = 4.80 (OL) – 2.50 $r^2 = 0.869$
Weight: WT = 0.0238 SL$^{2.692}$ $r^2 = 0.932$



Hippoglossoides elassodon
(flathead sole)

Left otolith; length (mm): 5.5
Fish length (cm): 19.1
Fish weight (g): 117.3

Regression equations:

Length: FL = 4.63 (OL) – 0.71 $r^2 = 0.947$
Weight: WT = 0.0078 FL$^{3.041}$ $r^2 = 0.984$



Hippoglossus stenolepis
(Pacific halibut)

Left otolith; length (mm): 12.8
Fish length (cm): 80.0
Fish weight (g): 6,600.0

Regression equations:

Length: No data available
Weight: No data available



Hypsopsetta guttulata
(diamond turbot)

Left otolith; length (mm): 4.8
Fish length (cm): 21.3
Fish weight (g): 345.0

Regression equations:

Length: SL = 4.89 (OL) – 0.29 $r^2 = 0.835$
Weight: WT = 0.0853 SL$^{2.664}$ $r^2 = 0.967$

PLEURONECTIDAE (cont.)



Hypsopsetta guttulata
(diamond turbot)

Right otolith; length (mm): 4.7
Fish length (cm): 21.3
Fish weight (g): 345.0

Regression equations:

Length: SL = 4.89 (OL) – 0.29 $r^2 = 0.835$
Weight: WT = 0.0853 SL$^{2.664}$ $r^2 = 0.967$



Microstomus pacificus
(Dover sole)

Left otolith; length (mm): 5.3
Fish length (cm): 26.9
Fish weight (g): 268.1

Regression equations:

Length: SL = 3.72 (OL) + 6.97 $r^2 = 0.587$
Weight: WT = 0.0094 SL$^{3.092}$ $r^2 = 0.854$



Microstomus pacificus
(Dover sole)

Right otolith; length (mm): 5.4
Fish length (cm): 26.9
Fish weight (g): 268.1

Regression equations:

Length: SL = 3.72 (OL) + 6.97 $r^2 = 0.587$
Weight: WT = 0.0094 SL$^{3.092}$ $r^2 = 0.854$

Microstomus pacificus
(Dover sole)

Right otolith; length (mm): 4.6
Fish length (cm): 26.2
Fish weight (g): 239.5

Regression equations:

Length: SL = 3.72 (OL) + 6.97 $r^2 = 0.587$
Weight: WT = 0.0094 SL$^{3.092}$ $r^2 = 0.854$

PLEURONECTIDAE (cont.)



Microstomus pacificus
(Dover sole)

Left otolith; length (mm): 4.8
Fish length (cm): 26.7
Fish weight (g): 289.0

Regression equations:

Length: SL = 3.72 (OL) + 6.97 $r^2 = 0.587$
Weight: WT = 0.0094 $SL^{3.092}$ $r^2 = 0.854$



Platichthys stellatus
(starry flounder)

Left otolith; length (mm): 7.0
Fish length (cm): 27.6
Fish weight (g): 547.4

Regression equations:

Length: SL = 3.35 (OL) + 0.23 $r^2 = 0.814$
Weight: WT = 0.0107 $SL^{3.268}$ $r^2 = 0.985$



Pleuronectes asper
(yellowfin sole)

Left otolith; length (mm): 6.6
Fish length (cm): 25.3
Fish weight (g): 294.4

Regression equations:

Length: SL = 2.17 (OL) + 10.65 $r^2 = 0.638$
Weight: WT = 0.0024 $SL^{3.605}$ $r^2 = 0.859$



Pleuronectes bilineatus
(rock sole)

Left otolith; length (mm): 6.8
Fish length (cm): 23.5
Fish weight (g): 287.2

Regression equations:

Length: FL = 6.16 (OL) – 6.97 $r^2 = 0.841$
Weight: WT = 0.0112 $FL^{2.997}$ $r^2 = 0.931$

PLEURONECTIDAE (cont.)



Pleuronectes vetulus
(English sole)

Right otolith; length (mm): 6.6
Fish length (cm): 21.2
Fish weight (g): 104.0

Regression equations:

Length: SL = 3.82 (OL) – 2.76 $r^2 = 0.965$
Weight: WT = 0.0163 $SL^{2.939}$ $r^2 = 0.995$



Pleuronectes vetulus
(English sole)

Left otolith; length (mm): 9.0
Fish length (cm): 32.3
Fish weight (g): 439.2

Regression equations:

Length: SL = 3.82 (OL) – 2.76 $r^2 = 0.965$
Weight: WT = 0.0163 $SL^{2.939}$ $r^2 = 0.995$



Psettichthys melanostictus
(sand sole)

Right otolith; length (mm): 5.2
Fish length (cm): 23.0
Fish weight (g): 288.0

Regression equations:

Length: SL = 5.06 (OL) – 3.18 $r^2 = 0.942$
Weight: WT = 0.0052 $SL^{3.441}$ $r^2 = 0.983$



Psettichthys melanostictus
(sand sole)

Left otolith; length (mm): 4.5
Fish length (cm): 21.7
Fish weight (g): 231.3

Regression equations:

Length: SL = 5.06 (OL) – 3.18 $r^2 = 0.942$
Weight: WT = 0.0052 $SL^{3.441}$ $r^2 = 0.983$

SOLEIDAE



Symphurus atricauda
(California tonguefish)

Left otolith; length (mm): 2.3
Fish length (cm): 14.5
Fish weight (g): 33.2

Regression equations:

Length: SL = 3.56 (OL) + 4.64 $r^2 = 0.464$
Weight: WT = 0.0074SL$^{3.136}$ $r^2 = 0.789$

NOAA Technical Reports NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for their optimum use. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, with the development and enforcement of domestic fisheries regulations, with the surveillance of foreign fishing off U.S. coastal waters, and with the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing services and economic analysis programs and through mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

Recently Published Technical Reports

138. Southeastern U.S. deepwater reef fish assemblages, habitat characteristics, catches, and life history summaries, by R. O. Parker Jr. and R. W. Mays. September 1998, 41 p.

139. Seasonal, horizontal, and vertical distribution of phytoplankton chlorophyll *a* in the northeast U.S. continental shelf ecosystem, by John E. O'Reilly and Christine Zetlin. November 1998, 120 p.

140. Quantitative composition and distribution of the macrobenthic invertebrate fauna of the continental shelf ecosystems of the northeastern United States, by Roger B. Theroux and Roland L. Wigley. December 1998, 240 p.

141. Marine flora and fauna of the eastern United States. Anthozoa: Actiniaria, Corallimorpharia, Ceriantharia, and Zoanthidea, by Kenneth P. Sebens. December 1998, 68 p.

142. Biology and fisheries of swordfish, *Xiphias gladius*. Papers from the International Symposium on Pacific Swordfish, Ensenada, Mexico, 11-14 December 1994, edited by Izadore Barrett, Oscar Sosa-Nishizaki, and Norman Bartoo. December 1998, 276 p.

143. Characteristics and causes of Texas marine strandings, edited by Roger Zimmerman. December 1998, 85 p.

144. Distribution of Atlantic menhaden, *Brevoortia tyrannus*, purse-seine sets and catches from southern New England to North Carolina, 1985-96, by Joseph W. Smith. March 1999, 22 p.

145. Distribution patterns of fish eggs in the U.S. northeast continental shelf ecosystem, 1977-1987, by Peter Berrien and John Sibunka. June 1999, 310 p.

146. Synopsis of biological data on the Nassau grouper, *Epinephelus striatus* (Bloch, 1792), and the jewfish, *E. itajara* (Lichtenstein, 1822), by Yvonne Sadovy and Anne-Marie Eklund. July 1999, 65 p.

147. Investigation of unusual mortalities of bottlenose dolphins along the mid-Texas coastal bay ecosystem during 1992, by A. A. Colbert, G. I. Scott, M. H. Fulton, E. F. Wirth, J. W. Daugomah, P. B. Key, E. D. Strozier, and S. B. Galloway. November 1999, 23 p.

148. The distribution and size composition of finfish, American lobster, and long-finned squid in Long Island Sound based on the Connecticut Fisheries Division Bottom Trawl Survey, 1984-1994, by Kurt F. Gottschall, Mark W. Johnson, and David G. Simpson. February 2000, 195 p.

149. Population characteristics of Gulf menhaden, *Brevoortia patronus*, by Douglas S. Vaughan, Joseph W. Smith, and Michael H. Prager. April 2000, 19 p.

NOAA Technical Reports NMFS

Technical Reports of the *Fishery Bulletin*

Guide for Contributors

Preparation

Title page should include authors' full names and mailing addresses and the senior author's telephone and FAX number.

Abstract should not exceed one double-spaced typed page. It should state the main scope of the research but emphasize its conclusions and relevant findings. Because abstracts are circulated by abstracting agencies, it is important that they represent the research clearly and concisely.

Text must be typed double-spaced throughout. A brief introduction should portray the broad significance of the paper; the remainder of the paper should be divided into the following sections: **Materials and methods, Results, Discussion** (or **Conclusions**), and **Acknowledgments**. Headings within each section must be short, reflect a logical sequence, and follow the rules of multiple subdivision (i.e. there can be no subdivision without at least two items). The entire text should be intelligible to interdisciplinary readers; therefore, all acronyms, abbreviations, and technical terms should be spelled out the first time they are mentioned. The scientific names of species must be written out the first time they are mentioned; subsequent mention of scientific names may be abbreviated. Follow the *U.S. Government Printing Office Style Manual* (1984 ed.) and the *CBE Style Manual* (5th ed.) for editorial style, and the most current issue of the *American Fisheries Society's Common and Scientific Names of Fishes from the United States and Canada* for fish nomenclature. Dates should be written as follows: 11 November 1991. Measurements should be expressed in metric units, e.g., metric tons as (t); if other units of measurement are used, please make this fact explicit to the reader. The numeral one (1) should be typed as a one, not as a lower-case el (l).

Text footnotes should be numbered with Arabic numerals and typed on a separate sheet of paper. Footnote all personal communications, unpublished data, and unpublished manuscripts with full address of the communicator or author, or, as in the case of unpublished data, where the data are on file. Authors are advised to avoid references to nonstandard (gray) literature, such as internal, project, processed, or administrative reports, wherever possible. Where these references are used, please include whether they are available from NTIS (National Technical Information Service) or from some other public depository.

Literature cited comprises published works and those accepted for publication in peer-reviewed literature (in press). Follow the name and year system for citation format. In the text, cite Smith and Jones (1977) or (Smith and Jones, 1977). If there is a sequence of citations, list chronologically: Smith, 1932; Green, 1947; Smith and Jones, 1985. Abbreviations of serials should conform to abbreviations given in *Serial Sources for the BIOSIS Previews Database*. Authors are responsible for the accuracy and completeness of all citations.

Tables should not be excessive in size and must be cited in numerical order in the text. Headings should be short but ample enough to allow the table to be intelligible on its own. All unusual symbols must be explained in the table heading. Other incidental comments may be footnoted with italic numerals. Use asterisks for probability in statistical data. Because tables are typeset, they need only be submitted typed and formatted, with double-spaced legends. Zeros should precede all decimal points for values less than one.

Figures include line illustrations and photographs (or slides) and must be cited in numerical order in the text. Unless photographs are submitted on glossy paper with good contrast, we cannot guarantee a good final printed copy. Figures are to be labeled with author's name and number of figure. Use Times Roman font (upper and lowercase letters) to label within figures. Avoid vertical lettering except for *y*-axis labels. Zeros should precede all decimal points for values less than one. Figures may be submitted as computer software files (along with laser-printed copies), as photo-mechanical transfers (PMTs), or as high quality photographic prints. Send only xerox copies of figures to the Scientific Editor; original figures will be requested later when the manuscript has been accepted for publication. Figure legends should explain all symbols and abbreviations and should be double-spaced on a separate page at the end of the manuscript.

Copyright law does not cover government publications; they fall within the public domain. If an author reproduces any part of a government publication in his work, reference to source is considered correct form.

Submission

Send printed copies (original and three copies) to the Scientific Editor:

Dr. John V. Merriner, Scientific Editor
Southeast Fisheries Science Center
National Marine Fisheries Service, NOAA
101 Pivers Island Road
Beaufort, NC 02516

Once the manuscript has been accepted for publication, you will be asked to submit a software copy of your manuscript to the Managing Editor. The software copy should be submitted in WordPerfect or Word format and should be placed on a 3.5-inch disk that is double-sided, double or high density, and that is compatible with DOS/Windows systems. Note: We are no longer able to accept ASCII text files.

Copies of published reports (and articles within multi-authored reports) are available free of charge to the senior author (50 copies) and to his or her laboratory (50 copies). Additional copies may be purchased in lots of 100 when the author receives page proofs.

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC
ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS OFFICE
BIN C15700
SEATTLE, WA 98115
OFFICIAL BUSINESS

Penalty for Private Use, $300

BULK RATE
POSTAGE & FEES PAID
U.S. Department of Commerce
Permit No. G-19

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 13, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS—Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS—Reports containing data, observations, instructions, etc. A partial listing includes data serials; predictions and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

Information on availability of NOAA publications can be obtained from:

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161

☆ U. S. GOVERNMENT PRINTING OFFICE: 2000 PROG 2057-S/800015